arXiv:1212.2530v1 [math.CO] 11 Dec 2012

Pattern Avoidance in Ordered Set Partitions

Anant Godbole[1]
Department of Mathematics and Statistics
East Tennessee State University
Johnson City, TN 37614, USA
`godbolea@etsu.edu`
Adam Goyt
Department of Mathematics
Minnesota State University
Moorhead, MN 56563 USA
`goytadam@mnstate.edu`
Jennifer Herdan[1]
Department of Mathematics and Statistics
Winona State University
Winona, MN 55987, USA
Lara Pudwell
Department of Mathematics and Computer Science
Valparaiso University
Valparaiso, IN 46383, USA
`Lara.Pudwell@valpo.edu`

Abstract

In this paper we consider the enumeration of ordered set partitions avoiding a permutation pattern of length 2 or 3. We provide an exact enumeration for certain special cases, and a recursive technique to exactly enumerate the appropriate set partitions in general. We also give some asymptotic results for the growth rates of the number of ordered set partitions avoiding a single pattern; including a Stanely-Wilf type result bounding such growth rates.

Keywords: ordered set partitions, pattern avoidance, permutations

1 Introduction

Pattern avoidance in permutations was first introduced by Knuth [11], and continues to be an active area of research today. Let $\mathcal{S}_n$ denote the set of permutations of length n, and consider $\pi \in \mathcal{S}_n$ and $\rho \in \mathcal{S}_m$. We say π *contains* ρ as a pattern if there exist indices

---

[1] Research of the first and third authors partially funded by NSF grant number DMS-1004624.

$1 \leq i_1 < i_2 < \cdots < i_{m-1} < i_m \leq n$ such that $\pi_{i_a} \leq \pi_{i_b}$ if and only if $\rho_a \leq \rho_b$. In this case, we say that $\pi_{i_1}\pi_{i_2}\cdots\pi_{i_m}$ is *order-isomorphic* to ρ. Otherwise, we say π *avoids* ρ. Further, let $\mathcal{S}_n(\rho)$ denote the set of permutations of length n that avoid ρ, and let $\mathrm{s}_n(\rho) = |\mathcal{S}_n(\rho)|$. It is straightforward to see that $\mathrm{s}_n(12) = 1$ for $n \geq 0$ because the only permutation of length n that avoids 12 is the decreasing permutation. It is also well-known that given any $\rho \in \mathcal{S}_3$, $\mathrm{s}_n(\rho) = C_n$ where $C_n = \dfrac{\binom{2n}{n}}{n+1}$ is the nth Catalan number [15].

Pattern avoidance has been studied in contexts other than permutations. In particular, the notion of pattern-avoidance in set partitions was introduced by Klazar [8], with further work done by Klazar, Goyt, and Sagan [3, 4, 9, 10, 14]. More recently, Goyt and Pudwell introduced the notion of colored set partitions and considered three distinct types of pattern avoidance in this context [5, 6]. In this paper, we consider a definition of pattern avoidance most closely related to that of [6].

A partition p of the set $S \subseteq \mathbb{Z}$, written $p \vdash S$, is a family of nonempty, pairwise disjoint subsets $B_1, B_2, \ldots, B_k$ of S called *blocks* such that $\cup_{i=1}^{k} B_i = S$. We write $p = B_1/B_2/\ldots/B_k$ and define the *length* of p, written $\ell(p)$, to be the number of blocks. Note that because $B_1, \ldots, B_k$ are sets, the order of elements within a block does not matter; for convenience we will write elements of a block in increasing order. We are particularly interested in the set of *ordered partitions of* $[n] = \{1, \ldots, n\}$ *into* k *blocks*, written $\mathcal{OP}_{n,k}$, which is the set of partitions p such that $p \vdash [n]$, $\ell(p) = k$, and where the order of blocks is important. For example 13/2/4 and 4/13/2 are two distinct members of $\mathcal{OP}_{4,3}$. In the sequel, we also let $\mathcal{OP}_n = \cup_{k=1}^{n} \mathcal{OP}_{n,k}$, and let $\mathcal{OP}_{[b_1,\ldots,b_k]}$ be the set of ordered partitions p such that $p \vdash [b_1 + \cdots + b_k]$, $\ell(p) = k$ and $|B_i| = b_i$ for $1 \leq i \leq k$. Similarly, we let $\mathrm{op}_{n,k} = |\mathcal{OP}_{n,k}|$, $\mathrm{op}_n = |\mathcal{OP}_n|$, and $\mathrm{op}_{[b_1,\ldots,b_k]} = \left|\mathcal{OP}_{[b_1,\ldots,b_k]}\right|$

Given a partition $p \vdash [n]$, and a permutation $\rho \in \mathcal{S}_m$, we say that p contains ρ if there exist blocks $B_{i_1}, \ldots, B_{i_m}$ where $i_1 < i_2 < \cdots < i_m$, and there exists $b_j \in B_{i_j}$ such that $b_1 \cdots b_m$ is order-isomorphic to ρ. For example, $14/56/2/3 \in \mathcal{OP}_{6,4}$ contains the pattern $\rho = 312$, as evidenced by $b_1 = 4$, $b_2 = 2$, and $b_3 = 3$.

Avoidance in ordered partitions is attractive in that three special cases are directly related to other known enumerative results. First, note that for any $\rho \in \mathcal{S}_m$, we have $\mathrm{s}_n(\rho) = \mathrm{op}_{[\underbrace{1,\ldots,1}_{n}]}(\rho)$. That is, a permutation is equivalent to an ordered partition where all blocks are of size 1. Second, we will see in the next section that $\mathcal{OP}_{n,k}(12)$ is in bijection with the set of integer compositions of n into k parts. Finally, the definition of avoidance for ordered partitions described above corresponds to the pattern-type avoidance detailed in [6] if we considered unordered partitions where all elements in block i are given color c_i, and avoid the colored pattern $\rho_1^1\rho_2^2\cdots\rho_m^m$ rather than pattern ρ.

In Section 2, we give closed formulas for $\mathrm{op}_{n,k}(12)$, $\mathrm{op}_{n,3}(123)$, $\mathrm{op}_{n,3}(132)$, and $\mathrm{op}_{n,n-1}(132)$. In Section 3, we give a bijective proof that $\mathrm{op}_{[b_1,\ldots,b_k]}(123) = \mathrm{op}_{[b_1,\ldots,b_k]}(132)$ for any list of positive integers $b_1, \ldots, b_k$, settling the question of whether $\mathrm{op}_{[b_1,\ldots,b_k]}(\rho_1) = \mathrm{op}_{[b_1,\ldots,b_k]}(\rho_2)$ for any patterns $\rho_1, \rho_2 \in \mathcal{S}_3$. In Section 4, we adapt the enumeration schemes of Zeilberger [21], Vatter [19], and Pudwell [13] to enumerate the set $\mathcal{OP}_{[b_1,\ldots,b_k]}(123)$ for any block sizes $b_1, \ldots, b_k$. In Section 5 we investigate the case of $\mathrm{op}_{[2,2,\ldots 2]}(123)$ and continue with an analysis of $\mathrm{op}_{n,n-1}(123)$ in Section 6, where we also prove a monotonicity result. Then, in Section 7, we prove that a Stanley-Wilf limit exists as n tends to infinity for all the sequences $\mathrm{op}_{n,k}(\rho)$ where k and ρ are held fixed. We end with some open questions.

2 A few simple cases

In this section, we consider a few special cases of the pattern avoidance problem introduced in the introduction. In particular, we consider $\mathrm{op}_{n,k}(12)$, $\mathrm{op}_{n,3}(123)$, $\mathrm{op}_{n,3}(132)$, and $\mathrm{op}_{n,n-1}(132)$. The simplest of these cases is that of avoiding the pattern $\rho = 12$.

Theorem 1.

$$\mathrm{op}_{n,k}(12) = \binom{n-1}{k-1}$$

Proof. Notice that a member of $\mathcal{OP}_{n,k}$ avoids the pattern $\rho = 12$ if and only if for each block B_i, all elements of each block B_j where $j > i$ are strictly less than all elements of block B_i. Once we know the sizes of blocks $B_1, B_2, \ldots, B_k$, the 12-avoiding partition is determined: the largest $|B_1|$ elements are in block 1, the next largest $|B_2|$ elements are in block 2, and so on. Thus, $\mathrm{op}_{n,k}(12)$ is merely the number of integer compositions of n into k parts, which is well known to be $\binom{n-1}{k-1}$. □

Note that a similar argument shows that $\mathrm{op}_{n,k}(21) = \binom{n-1}{k-1}$. In fact, as with pattern-avoiding permutations, a few natural symmetries simplify our work. Given permutation pattern $\rho = \rho_1 \cdots \rho_m$, define the *reversal* of ρ, written ρ^r, to be $\rho^r = \rho_m\rho_{m-1}\cdots\rho_2\rho_1$, and define the *complement* of ρ, written ρ^c, to be $\rho^c = (m+1-\rho_1)(m+1-\rho_2)\cdots(m+1-\rho_m)$. Since complement and reversal both provide involutions on the set of permutations and on the set $\mathcal{OP}_{n,k}$, we have that $\mathrm{op}_{n,k}(\rho) = \mathrm{op}_{n,k}(\rho^r) = \mathrm{op}_{n,k}(\rho^c) = \mathrm{op}_{n,k}((\rho^r)^c)$. Unlike the case of pattern-avoiding permutations, we no longer have a well-defined notion of inverse, so reversal and complementation are the only natural symmetries of which we may take advantage.

For the case of patterns of length 3, we see that $123^r = 321$, $132^r = 231$, $231^c = 213$, and $213^r = 312$, so we have that $\mathrm{op}_{n,k}(123) = \mathrm{op}_{n,k}(321)$ and $\mathrm{op}_{n,k}(132) = \mathrm{op}_{n,k}(231) = \mathrm{op}_{n,k}(213) = \mathrm{op}_{n,k}(312)$. In Section 3 we will show that $\mathrm{op}_{n,k}(123) = \mathrm{op}_{n,k}(132)$ by demonstrating the stronger claim that $\mathrm{op}_{[b_1,\ldots,b_k]}(123) = \mathrm{op}_{[b_1,\ldots,b_k]}(132)$ for any choice of block sizes $b_1, b_2, \ldots, b_k$. For now, though, we consider the special case where $k = 3$.

Theorem 2.

$$\mathrm{op}_{n,3}(123) = \left(\frac{n^2}{8} + \frac{3n}{8} - 2\right)2^n + 3$$

Proof. It is easy to see that

$$\mathrm{op}_{n,3}(123) = \sum_{a=1}^{n-2}\sum_{b=1}^{n-1-a}\sum_{\ell=1}^{n-a+1}\sum_{i=\max\{0,\ell-1-c\}}^{\min\{\ell-1,b\}}\binom{\ell-1}{i}\binom{n-\ell}{a-1},$$

where $a, b, c = n - a - b$ represent the sizes of blocks 1, 2, and 3 respectively, ℓ is the smallest number in block 1, and i represents how many numbers smaller than ℓ are in block 2. Using standard algebraic techniques the above sum above may be reduced to $\left(\frac{n^2}{8} + \frac{3n}{8} - 2\right) 2^n + 3$. □

Next, we consider $\mathrm{op}_{n,3}(132)$. We will show why $\mathrm{op}_{n,3}(123) = \mathrm{op}_{n,3}(132)$ via a bijection in the next section, but first, we consider how to count members of $\mathcal{OP}_{n,3}(132)$ directly.

Theorem 3.

$$\mathrm{op}_{n,3}(132) = \left(\frac{n^2}{8} + \frac{3n}{8} - 2\right) 2^n + 3$$

Proof. Consider $p \in \mathcal{OP}_{n,3}(132)$. We have two cases: either p avoids the pattern 12 in the first two blocks, or p contains the pattern 12 in the first two blocks.

In the first case, there is no restriction on the values of elements in the third block, so we may choose a elements $(1 \leq a \leq n-2)$ to be members of B_3. Since we assume that there is no 12 pattern in the first two blocks, once we choose the number b $(1 \leq b \leq n - a - 1)$ of elements of B_2, we know the smallest b elements of B_1 and B_2 are in B_2, and the remaining elements are in B_1. Thus, there are $\sum_{a=1}^{n-2} \sum_{b=1}^{n-a-1} \binom{n}{a}$ possible partitions of this form.

The second case is more complicated. Since we know there is a 12 pattern in the first two blocks, let i be the smallest element of B_1 that participates in a 12 pattern and let j be the largest element of B_2 that participates in a 12 pattern. By definition of i and j, we know that there are no elements of B_1 smaller than i and there are no elements of B_2 larger than j. Further, there are no elements of B_3 that are both greater than i and less than j. Let a_1 be the number of elements of B_2 that are greater than i but less than j, and let a_2 be the number of elements of B_2 that are less than i. Further, let b be the number of elements of B_1 that are greater than j. Once we have determined, i, j, a_1, a_2, and b, it remains to choose which $a_1 + a_2$ elements appear in B_2 and which b elements appear in B_1. This yields $\sum_{i=1}^{n-1} \sum_{j=i+1}^{n} \sum_{a_1=0}^{j-i-1} \sum_{a_2=0}^{i-1} \sum_{b=0}^{n-j} \binom{j-i-1}{a_1} \binom{i-1}{a_2} \binom{n-j}{b} - \sum_{i=1}^{n-1} \sum_{j=i+1}^{n} \sum_{a1=0}^{j-i-1} \binom{j-i-1}{a_1}$ possible ordered partitions.

Together, we have that

$$\begin{aligned}
\mathrm{op}_{n,3}(132) = &\sum_{a=1}^{n-2} \sum_{b=1}^{n-a-1} \binom{n}{a} \\
&+ \sum_{i=1}^{n-1} \sum_{j=i+1}^{n} \sum_{a_1=0}^{j-i-1} \sum_{a_2=0}^{i-1} \sum_{b=0}^{n-j} \binom{j-i-1}{a_1} \binom{i-1}{a_2} \binom{n-j}{b} \\
&- \sum_{i=1}^{n-1} \sum_{j=i+1}^{n} \sum_{a_1=0}^{j-i-1} \binom{j-i-1}{a_1},
\end{aligned}$$

and through standard algebraic manipulation, these sums simplify to

$$\mathrm{op}_{n,3}(132) = \left(\frac{n^2}{8} + \frac{3n}{8} - 2\right) 2^n + 3,$$

as desired. □

Note that having $k = 3$ blocks is the minimum non-trivial number of blocks to consider when avoiding a pattern of length 3. Similarly, as mentioned above, $k = n$ blocks is equivalent to considering pattern-avoiding permutations, so the maximum non-trivial number of blocks to consider is $k = n - 1$. We conclude this section with a an enumeration result in this maximal-block case.

Let $\rho = \rho_1\rho_2 \dots \rho_n \in \mathcal{S}_n$. We say ρ_i is a *right-to-left maximum* if $\rho_i > \rho_j$ for any $j > i$. Let $r(n, k)$ be the number of $\rho \in \mathcal{S}_n$ with k right-to-left maxima. Note that the last element of a permutation is always a right-to-left maximum.

Lemma 1. *Let $\rho \in \mathcal{S}_n(132)$, and suppose $m > n$. The only position where m could be placed in ρ without introducing a copy of 132 is immediately after a right-to-left maximum or at the beginning of ρ.*

Proof. Let $\rho = \rho_1\rho_2 \dots \rho_n \in \mathcal{S}_n(132)$ and $m > n$. Suppose we want to insert m at the beginning, the end or between ρ_i and ρ_{i+1} and not introduce a new copy of 132. A copy of 132 involving m must have m representing 3. Thus, m could have been placed at the beginning or end of ρ. Suppose m is not placed at the beginning or end of ρ, and a copy of 132 is formed. Looking to the left of m find the first right-to-left maximum or ρ, call it ρ_k. Since ρ avoids 132 the element of ρ representing the 1 in this copy of 132 must appear between ρ_k and m. Thus, placing m immediately to the right of a right-to-left maximum will produce a permutation that avoids 132. We will show that this is the only place that on could place m.

Suppose that we do not place m immediately to the right of a right-to-left maximum or at the beginning or end. Suppose again that ρ_k is the first right-to-left maximum appearing to the left of m. There must be an element between ρ_k and m call it j. Now j cannot be a right-to-left maximum, and m is not at the end of ρ so there must be a right-to-left maximum appearing after m. Together j, m and this right-to-left maximum form a copy of 132. Thus, the only place that m could be placed is immediately to the right of a right-to-left maximum or at the beginning of ρ. □

Theorem 4.

$$\begin{aligned}
\mathrm{op}_{n,n-1}(132) &= \sum_{k=0}^{n-2} C_k + \sum_{a=2}^{n-1} \sum_{k=0}^{n-a-2} C_k \sum_{i=1}^{a-1} r(a-1, i) \binom{i+n-k-a-1}{i} \\
&+ \sum_{a=2}^{n-1} C_{n-a-1} C_{a-1} + \sum_{a=2}^{n-1} \sum_{k=n-a}^{n-2} C_{n-a-1} C_{k-n+a+1} C_{n-k-2} \\
&+ 2 \sum_{j=1}^{n} C_{j-1} \, \mathrm{op}_{n-j,n-j-1}(132).
\end{aligned}$$

Proof. Let $\rho \in \mathcal{OP}_{n,n-1}(132)$. Since ρ has $n-1$ blocks there must be exactly one block of size two (doubleton) and the remaining blocks are of size one. We will focus on this doubleton.

Suppose first of all that n is not in the doubleton. Thus, n is in its own block and we condition on the location of n. If n is in the j^{th} position then each of the $j-1$ elements to the left of n must be larger than each of the $n-j$ elements to the right of n else n is the 3 in a 132 pattern. Either the doubleton appears to the right or to the left of n. If it appears to the right of n then there are $\mathrm{op}_{n-j,n-j-1}(132)$ partitions of the $n-j$ elements to the right of n with $n-j-1$ blocks that avoid 132, and as is well known there are C_{j-1} partitions of the $j-1$ elements to the left of n with $j-1$ blocks which avoid 132. So there are $\sum_{j=1}^{n} C_{j-1}\mathrm{op}_{n-j,n-j-1}(132)$ ordered partitions of $[n]$ with $n-1$ blocks avoiding 132 with the doubleton not containing n and the doubleton appearing to the right of n. The same sum counts those with the doubleton appearing to the left of n. This gives us the last term in the sum.

We now consider the partitions with n in the doubleton. We will condition on the element, a, joining n in the doubleton. First suppose that $a = 1$. Suppose that there are k elements to the left of the doubleton. Since n is in the doubleton, these k elements must be larger than the $n-k-2$ elements that appear to the right of the doubleton. Since the doubleton is $\{1, n\}$, every element to the right of the doubleton must be in increasing order. The k elements to the left of the doubleton must simply avoid 132, and hence there are C_k partitions of these k elements that avoid 132. Thus, there are $\sum_{k=0}^{n-2} C_k$ partitions of n with $n-1$ blocks avoiding 132 where the doubleton is $\{1, n\}$.

Now, suppose that $2 \le a \le n-1$. Observe that if there are k elements to the left of $\{a, n\}$ then these must be the k largest remaining elements, and the $n-k-2$ smallest remaining elements appear to the right of $\{a, n\}$. If the k elements to the left of $\{a, n\}$ are all larger than a then there can be no copy of 132 involving any of these and a. There are $n-a-1$ elements that are larger than a. So if $k \le n-a-1$ then there are C_k ways to arrange the elements that appear to the left of $\{a, n\}$. If $k \ge n-a$ then of these k elements the $n-a-1$ that are larger than a must appear before the $k-n+a+1$ that are smaller than a. There are $C_{n-a-1}C_{k-n+a+1}$ ways to arrange these elements. This gives us the first factor in the second and third terms of the recursion and the first factors of the fourth term of the recursion.

We need to consider what happens with the $n-k-2$ elements that appear after $\{a, n\}$. Assume that $k \le n-a-2$. This means that there is at least one element larger than a appearing after $\{a, n\}$. The elements that are larger than a appearing the right of a must be in increasing order, otherwise a copy of 132 is created. The $a-1$ smallest elements could only possibly form a copy of 132 with each other or with the $n-k-a-1$ elements appearing to the right of $\{a, n\}$ that are larger than a. We begin by forming a 132-avoiding permutation of the $a-1$ elements. Now, by Lemma 1, when placing the $n-k-a-1$ larger elements we may only place them immediately to the right of a right-to-left maximum or at the beginning of the permutation. There are $r(a-1, i)$ permutations of $[a-1]$ with i right-to-left maxima. Once we have selected such a permutation, ρ, with i right-to-left maxima there are $\binom{n-k-a-1+i}{i}$ ways to place the $n-k-a-1$ elements that are larger than a into the positions immediately following the right-to-left maxima of ρ, since these elements must be in increasing order. If we sum for $1 \le i \le a-1$, multiply by C_k from above and sum over appropriate choices for a and k we get the second term.

Now, suppose that $k = n - a - 1$. Then the remaining $a - 1$ elements are to the right of $\{a, n\}$. Thus, there are no elements larger than a on the right of $\{a, n\}$, so no 132 can be formed involving a. So there are $C_{n-a-1}C_{a-1}$ partitions with $n - a - 1$ elements to the left of $\{a, n\}$. This is the third term.

Finally, if $k \geq n - a$. Again all of the elements to the right of $\{a, n\}$ are smaller than a, so there are C_{n-k-2} arrangements of these elements that avoid 132. Multiplying by $C_{n-a-1}C_{k-n+a+1}$ as described above and summing over appropriate values of a and k gives us the fourth term in the recursion. $\square$

Of course, it would be nice to know the value of $r(n, k)$ so that we have a nice recursive formula for computing $\mathrm{op}_{n,n-1}(132)$. It turns out that $r(n, k)$ is closely related to the Catalan numbers.

Theorem 5. *Let $r(n, k)$ be the number of permutations in $S_n(132)$ that have k right-to-left maxima. Then $r(0, 0) = 1$, $r(n, 0) = 0$ if $n > 0$ and $r(n, k) = 0$ if $k > n$. Furthermore, let $C_n^{(k)} = [x^n](C(x))^k$, then for $n \geq k$ we have $r(n, k) = C_{n-k}^{(k)}$.*

Proof. We define $r(0, 0) = 1$ by considering the empty partition. If $n \geq 1$ the last element in a permutation in $S_n(132)$ is always a right-to-left maximum, so there must be at least one right-to-left maximum. Hence $r(n, 0) = 0$ if $n \geq 1$. Also, there cannot be more right-to-left maxima than there are entries in the permutation. Thus, $r(n, k) = 0$ if $k > n$.

Let $\rho \in S_n(132)$ have k right-to-left maxima. Now, suppose that n is in the j^{th} position. Then the first $j - 1$ positions must consist of the $j - 1$ largest elements of $[n]$ and avoid 132. Furthermore, since these elements precede n, none of them could be a right-to-left maximum. Thus, there are C_{j-1} ways to arrange the first $j - 1$ elements.

The last $n - j$ elements of ρ must be the $n - j$ smallest elements of $[n]$, avoid 132 and consist of $k - 1$ right-to-left maxima. There are $r(n - j, k - 1)$ of these. Thus, we have that $r(n, k) = \sum_{j=1}^{n} C_{j-1}r(n - j, k - 1)$, for $n \geq k$ and $k \geq 1$.

This recursive form gives us that $r(n, 1) = C_{n-1}$ for $n \geq 1$. Thus, we have that $r(n, 2) = \sum_{j=1}^{n} C_{j-1}r(n - j, 2 - 1) = \sum_{j=1}^{n-1} C_{j-1}C_{n-j-1} = C_{n-2}^{(2)}$ for $n \geq 2$. By induction we have that $r(n, k) = C_{n-k}^{(k)}$ for $n \geq k$. $\square$

Callan [2] proves that $C_n^{(k)} = \frac{k}{2n+k}\binom{2n+k}{n}$. This gives us that $r(n, k) = \frac{k}{n}\binom{2n-k-1}{n-k}$. Thus, we restate Theorem 4.

Theorem 4.

$$\begin{aligned} \mathrm{op}_{n,n-1}(132) \quad = \quad & \sum_{k=0}^{n-2} C_k + \sum_{a=2}^{n-1} \sum_{k=0}^{n-a-2} C_k \sum_{i=1}^{a-1} \frac{i}{a-1}\binom{2a+k-3}{a-2}\binom{i+n-k-a-1}{i} + \\ & \sum_{a=2}^{n-1} C_{n-a-1}C_{a-1} + \sum_{a=2}^{n-1} \sum_{k=n-a}^{n-2} C_{n-a-1}C_{k-n+a+1}C_{n-k-2} + \\ & 2\sum_{j=1}^{n} C_{j-1}\,\mathrm{op}_{n-j,n-j-1}(132). \end{aligned}$$

3 A Bijection and Pattern Avoidance in Words

As was mentioned in the previous section, the usual symmetries of reversal and complementation are not enough to show that $\mathrm{op}_{[b_1,\ldots,b_k]}(123) = \mathrm{op}_{[b_1,\ldots,b_k]}(132)$. We will do this by adapting the familiar bijection of Simion and Schmidt [11]. To use their bijection we will need a notion of left-to-right minima for set partitions. Let $p = B_1/B_2/\ldots/B_k \in \mathcal{OP}$, then we will say that element $a \in B_i$ is a *left-to-right minimum* if a is smaller than every element appearing in blocks B_j for $1 \leq j \leq i-1$.

We will first describe the bijection through example. Consider the ordered partition 59/38/1267/4. This partition avoids 123. Notice that the left-to-right minima in this partition are 1, 2, 3, 5, and 9. Also, notice that the other elements in this partition form a decreasing sequence if we place the elements in the same block in decreasing order. Remove the elements that are not left-to-right minima. Now we have the partition $59/3/12/\emptyset$, where the second block is missing one element, the third is missing two elements and the last block is missing one element. We will fill the gaps in the blocks working from left to right by placing the smallest remaining elements that is larger than the smallest left-to-right minimum in the preceding block. So we would place the element 6 in block 2, obtaining $59/36/12/\emptyset$. We would then place 4 and 7 in the third block, producing the partition $59/36/1247/\emptyset$. Finally, we place 8 in the last block producing 59/36/1247/8. This partition is 132 avoiding.

The inverse of this bijection is achieved by placing all of the elements other than the left-to-right minima in descending order.

Theorem 6. *For $n \geq 1$, $\mathrm{op}_{[b_1,\ldots,b_k]}(123) = \mathrm{op}_{[b_1,\ldots,b_k]}(132)$.*

Proof. Let $\phi : \mathcal{OP}_{[b_1,\ldots,b_k]}(123) \to \mathcal{OP}_{[b_1,\ldots,b_k]}(132)$ be as follows. For $p \in \mathcal{OP}_{[b_1,\ldots,b_k]}(123)$, we will construct a corresponding partition $\phi(p) \in \mathcal{OP}_{[b_1,b_2,\ldots,b_k]}(132)$.

First find the left-to-right minima of p and leave them fixed. Now, remove the other elements of the partition. Working from left to right fill in the missing entries in each block by placing the smallest remaining element that is larger than the preceding left-to-right minimum. This new partition will avoid 132, since if a copy of 132 did appear, then one would appear with a left-to-right minimum representing the 1 in the copy of 132. This would imply that the element representing the 2 was placed in a block after then element representing the 3, which contradicts the prescribed placement of the elements. Thus, no copy of 132 appears.

The inverse of this construction is to again leave the left-to-right minima in place and place the remaining elements in descending order. Since the partition will essentially consist of two decreasing sequences, there is no way to form a copy of 123. $\square$

In his thesis [1], Burstein shows that the number of words avoiding the permutation 123 is the same as the number of words avoiding the permutation 132 using analytic techniques. Jelínek and Mansour [7] give a bijective proof of the same fact. It turns out that the bijection above can be used to give another bijective proof of this fact. We will need to first discuss how words and ordered partitions are related.

Using the concept of a permutation graph, we define an ordered partition graph to be a permutation graph where we allow more than one entry in a column. For example, the graph associated to the partition 4/13/256 is given in Figure 1.



Figure 1: Graph of 4/13/256

Let $k \geq 1$ and $n \geq 1$ then the set of words of length n with letters from the alphabet $[k]$ is denoted $[k]^n$. We say that a word $w \in [k]^n$ *contains* $u \in [\ell]^m$ if there are indices $i_1 < i_2 < \cdots < i_m$ such that the word $w_{i_1} w_{i_2} \ldots w_{i_m}$ is order isomorphic to u. Otherwise we say that w *avoids* u. We let $[k]^n(u)$ be the set of all $w \in [k]^n$ that avoid u. For example the word $232133 \in [3]^6$ avoids the word 123. Since permutations are words in $[n]^n$ without repeated letters, we can consider words that avoid permutations as well.

We may encode a word using a graph like that above. In the case of a word, however, we will allow more than one entry in a row. The word 232133 has graph.



Figure 2: Graph of 232133

As was mentioned in Section 2, we have three symmetries reversal and complementation which were defined, and inverse which is not a symmetry of ordered partitions, since the inverse operation will not create an ordered partition from any ordered partition with a block with more than one element. Since the inverse operation is applied to an ordered partition by simply reflecting the graph of the partition in the line $y = x$, we observe that an ordered partition becomes a word when we apply the inverse operation. Observe that the $232133^i = 4/13/256$, where i is the inverse operation.

This gives us that the inverse operation i is a bijection between the set of words $[k]^n$ and the set of ordered partitions with k blocks where we allow blocks to be empty. For example the word $255332255 \in [5]^9$ is mapped to the ordered partition $\emptyset/167/45/\emptyset/2389$, that is $255332255^i = \emptyset/167/45/\emptyset/2389$. Notice that the first and fourth blocks are empty. We also observe that the permutations 123 and 132 are fixed under the inversion operation. Thus, if a word w avoids 123 or 132 then so does w^i and similarly for ordered partitions.

Recall the map $\phi : \mathcal{OP}_{[b_1,\ldots,b_k]}(123) \to \mathcal{OP}_{[b_1,\ldots,b_k]}(132)$ from above. We may extend this map to ordered partitions where we allow some empty blocks, by simply leaving these blocks empty. For any $w \in [k]^n(123)$, we have that $(\phi(w^i))^i \in [k]^n(132)$. Let ψ be the operation where we invert a word to form an ordered partition then apply the map ϕ above and then invert again to produce a word. We note that ψ itself is an involution, and hence we have the following theorem:

Theorem 7. *For $n \geq 0$,*

$$|[k]^n(123)| = |[k]^n(132)|.$$

4 Enumeration schemes for ordered partitions

Now that we have investigated ordered partitions avoiding patterns of length 2 and 3, we consider a way to enumerate members of $\mathcal{OP}_{[b_1,\ldots,b_k]}(123)$ more generally. The ideas in this section are largely adapted from the notion of enumeration scheme. Informally, an enumeration scheme is a system of recurrences that enumerate members of a family of sets. Such enumeration schemes were first introduced by Zeilberger [21] in the context of pattern-avoiding permutations. Later, Vatter [19] improved the efficiency of the schemes and Pudwell [13] generalized enumeration schemes to apply to pattern-avoiding words, that is, permutations of multisets. One particularly attractive point of enumeration schemes in these contexts have been that the recurrences involved in enumeration schemes can be completely deduced by computer algorithm. Although such an algorithm has not yet been written for ordered partitions avoiding *any* permutation pattern, we capture many of the same ideas of Zeilberger, Vatter, and Pudwell and adapt them to the current context in order to find a family of recurrences that computes $\text{op}_{[b_1,\ldots,b_k]}(123)$ for any list $b_1, \ldots, b_k$ of block sizes.

First, we condition on the patterns formed by the members of the first few blocks of a partition. To this end, let

$$\mathcal{OP}_{[b_1,\ldots,b_k]}(123; s) = \{p \in \mathcal{OP}_{[b_1,\ldots,b_k]}(123) | B_1 = s\}$$

and $\text{op}_{[b_1,\ldots,b_k]}(123; s) = \left|\mathcal{OP}_{[b_1,\ldots,b_k]}(123; s)\right|$. For example, $\text{op}_{[2,1,1]}(123; \{2,4\}) = 2$ because $\mathcal{OP}_{[2,1,1]}(123; \{2,4\}) = \{24/1/3, 24/3/1\}$.

Similarly, let

$$\mathcal{OP}_{[b_1,\ldots,b_k]}(123; [s,t]) = \{p \in \mathcal{OP}_{[b_1,\ldots,b_k]}(123) | B_1 = s, B_2 = t\}$$

and $\text{op}_{[b_1,\ldots,b_k]}(123; [s,t]) = \left|\mathcal{OP}_{[b_1,\ldots,b_k]}(123; [s,t])\right|$. For example, $\text{op}_{[2,1,1]}(123; [\{2,4\},\{3\}]) = 1$ because $\mathcal{OP}_{[2,1,1]}(123; [\{2,4\},\{3\}]) = \{24/3/1\}$.

Notice that we immediately have $\text{op}_{[b_1]}(123; \{1,2,\ldots,b_1\}) = 1$ and $\text{op}_{[b_1,b_2]}(123; s) = 1$ for any set $s \subseteq [n]$ where $|s| = b_1$. More generally, we also see that

$$\mathcal{OP}_{[b_1,\ldots,b_k]}(123) = \bigcup_{s \subseteq [n], |s| = b_1} \mathcal{OP}_{[b_1,\ldots,b_k]}(123; s), \text{ and}$$

$$\mathcal{OP}_{[b_1,\ldots,b_k]}(123; s) = \bigcup_{t \subseteq [n] \setminus s, |t| = b_2} \mathcal{OP}_{[b_1,\ldots,b_k]}(123; [s,t]).$$

In addition, since the sets on the right hand side of each equation are disjoint, we have

$$\text{op}_{[b_1,\ldots,b_k]}(123) = \sum_{s \subseteq [n], |s| = b_1} \text{op}_{[b_1,\ldots,b_k]}(123; s), \text{ and} \tag{1}$$

$$\mathrm{op}_{[b_1,\ldots,b_k]}(123; s) = \sum_{t \subseteq [n]\setminus s, |t|=b_2} \mathrm{op}_{[b_1,\ldots,b_k]}(123; [s,t]). \tag{2}$$

We are interested in $\mathrm{op}_{[b_1,\ldots,b_k]}(123)$, which can be written in terms of $\mathrm{op}_{[b_1,\ldots,b_k]}(123; s)$. We have an exact value for $\mathrm{op}_{[b_1,\ldots,b_k]}(123; s)$ in a couple cases, but otherwise we write $\mathrm{op}_{[b_1,\ldots,b_k]}(123; s)$ in terms of $\mathrm{op}_{[b_1,\ldots,b_k]}(123; [s,t])$. It remains to find a recurrence for each $\mathrm{op}_{[b_1,\ldots,b_k]}(123; [s,t])$.

Notice that the sets s and t may interact in one of two ways: either all elements in s are larger than all elements of t, or they are not.

In the first case, when all elements of s are larger than all elements of t, we note that there is a bijection between the sets $\mathcal{OP}_{[b_1,\ldots,b_k]}(123; [s,t])$ and $\mathcal{OP}_{[b_2,\ldots,b_k]}(123; t)$. Consider $p \in \mathcal{OP}_{[b_1,\ldots,b_k]}(123; [s,t])$. Certainly, deleting the entire first block does not produce a new 123 pattern, so after replacing the ith smallest element of the remaining ordered partition with i, we have obtained a member of $\mathcal{OP}_{[b_2,\ldots,b_k]}(123; t)$. To show that this is indeed a bijection, we also show that no 123-containing partitions would map to $\mathcal{OP}_{[b_2,\ldots,b_k]}(123; t)$. To this end, suppose that p has block sizes $[b_1, \ldots, b_k]$, $B_1 = s$, $B_2 = t$, but p contains a 123 pattern. If this pattern does not involve elements from B_1, then deleting B_1 will still produce a partition that contains 123. If this pattern does involve an element from B_1, that element must play the role of "1" in the 123 pattern. However, since all elements of B_2 are less than all elements of B_1, the roles of "2" and "3" will be played by elements from $B_3, \ldots, B_k$. This means that even if B_1 were deleted, the resulting ordering partition will still contain 123, and we are done. Since there is a bijection between the sets $\mathcal{OP}_{[b_1,\ldots,b_k]}(123; [s,t])$ and $\mathcal{OP}_{[b_2,\ldots,b_k]}(123; t)$, we see that

$$\mathrm{op}_{[b_1,\ldots,b_k]}(123; [s,t]) = \mathrm{op}_{[b_2,\ldots,b_k]}(123; t). \tag{3}$$

The second case is more complicated. If not all elements of s are larger than all elements of t, then there exists a 12 pattern within the first two blocks. Let i be the smallest element of B_2 that plays the role of "2" in a 12 pattern. Further let a be the number of elements of B_1 that are less than i and let b be the number of elements of B_2 that are less than i. Because i plays the role of "2" in a 12 pattern, we know that there are no elements larger than i in blocks $B_3, \ldots, B_k$. This means that $\mathrm{op}_{[b_1,\ldots,b_k]}(123; [s,t]) = 0$ if $\min(s) < \max(t)$ and $([n] \setminus [a+b]) \not\subseteq s \cup t$.

Suppose then, that $\min(s) < \max(t)$ and $([n] \setminus [a+b]) \subseteq s \cup t$. In this case, i and all elements larger than it cannot be involved in a 123 pattern because these elements all appear in B_1 and B_2, and there are no elements of $B_3, \ldots, B_k$ that are larger than i. We have $\mathrm{op}_{[b_1,\ldots,b_k]}(123; [s,t]) = \mathrm{op}_{[a,b,b_3,\ldots,b_k]}(123, [s^*, t^*])$ where s^* is the smallest a elements of s and t^* is the smallest b elements of t. Further, since i was the smallest element of B_2 involved in a 12 pattern, we now know that all elements of s^* are larger than all elements of t^*, so by Equation 3, we see that $\mathrm{op}_{[a,b,b_3,\ldots,b_k]}(123, [s^*, t^*]) = \mathrm{op}_{[b,b_3,\ldots,b_k]}(123, t^*)$.

Together, we have that

$$\mathrm{op}_{[b_1,\ldots,b_k]}(123;[s,t]) = \begin{cases} \mathrm{op}_{[b_2,\ldots,b_k]}(123;t) & \min(s) > \max(t) \\ 0 & \min(s) < \max(t) \text{ and } ([n] \setminus [a+b]) \not\subseteq s \cup t \\ \mathrm{op}_{[b,b_3,\ldots,b_k]}(123,t^*) & \text{otherwise } (t^* \text{ is the } b \text{ least elements of } t). \end{cases} \tag{4}$$

Now, Equations 1, 2, 3 and 4 together with the base cases $\mathrm{op}_{[b_1]}(123;\{1,2,\ldots,b_1\}) = 1$ and $\mathrm{op}_{[b_1,b_2]}(123;s) = 1$ allow us to compute $\mathrm{op}_{[b_1,\ldots,b_k]}(123)$ for any list of block sizes. This family of recurrences has been implemented in the Maple package `123scheme` can be found on the fourth author's website `http://faculty.valpo.edu/lpudwell/maple.html`. Further, now that we have a way to compute $\mathrm{op}_{[b_1,\ldots,b_k]}(123)$, we may use this computation to find $\mathrm{op}_{n,k}(123)$ and $\mathrm{op}_n(123)$ for various values of n and k. Also, by the result of Section 3, these values will be the same for $\mathrm{op}_{[b_1,\ldots,b_k]}(\rho)$, $\mathrm{op}_{n,k}(\rho)$, and $\mathrm{op}_n(\rho)$ where ρ is any pattern of length 3.

In the next section, we use the data computed by our enumeration scheme to analyze the case of $\mathrm{op}_{[\underbrace{2,\ldots,2}_{k}]}(123)$.

5 The case of blocks of size 2

In this section, we consider the case where all partition blocks are of size two. We may, of course, use the techniques of Section 4 to compute $\mathrm{op}_{[\underbrace{2,\ldots,2}_{k}]}(123)$. The enumeration scheme provided in that section efficiently computes $\mathrm{op}_{[\underbrace{2,\ldots,2}_{k}]}(123)$, which is sequence A220097 in the Online Encyclopedia of Integer sequences [16]. However, further analysis with Zeilberger's `Maple` package *FindRec* [22] predicts that this sequence satisfies this second order linear recurrence:

$$\mathrm{op}_{[\underbrace{2,\ldots,2}_{k}]}(123) =$$

$$\frac{329k^3 - 749k^2 + 514k - 96}{2k(2k+1)(7k-9)}\mathrm{op}_{[\underbrace{2,\ldots,2}_{k-1}]}(123) + \frac{3(14k^3 - 39k^2 + 31k - 6)}{k(2k+1)(7k-9)}\mathrm{op}_{[\underbrace{2,\ldots,2}_{k-2}]}(123)$$

for $k > 3$, with $\mathrm{op}_{[2]}(123) = 1$ and $\mathrm{op}_{[2,2]}(123) = 6$. It remains an open problem to prove this recurrence describes our sequence in general.

The asymptotic behavior of this sequence resulting from this recurrence can be analyzed using the Birkhoff-Trjitzinsky method [20] (implemented in *AsyRec*), which predicts the behavior to be $\sim .1583 \cdot 12^k \dfrac{\left(1 - \frac{249}{392k}\right)}{k^{3/2}}$.

Since these experimental techniques predict the sequence $\mathrm{op}_{[\underbrace{2,\ldots,2}_{k}]}(123)$ to grow as 12^k, it is interesting to give an explicit analysis of this case to provide bounds on the asymptotics;

the lower bound of $\mathrm{op}^{1/k}_{[\underbrace{2,\ldots,2}_{k}]}(123) \geq 8$ can be exhibited by elementary means as follows: If π is a 123-avoiding permutation in S_{2k} (of which there are $\frac{\binom{4k}{2k}}{2k+1} \sim 16^k$), then π corresponds to an *ordered* 123-avoiding partition with k consecutive blocks of size 2 each, where the elements in each block are ordered. Each of these partitions corresponds to 2^k *ordered* partitions with part sizes 2, it follows that

$$2^k \mathrm{op}_{[\underbrace{2,\ldots,2}_{k}]}(123) \geq \frac{\binom{4k}{2k}}{2k+1} \sim 16^k,$$

which establishes the claim that $\mathrm{op}^{1/k}_{[\underbrace{2,\ldots,2}_{k/2}]}(123) \geq \sqrt{8}$.

6 The case of $n-1$ blocks and monotonicity of $\mathbf{op}_{n,k}(123)$

We may also use the techniques of Section 4 to compute $\mathrm{op}_{n,n-1}(123)$. The enumeration scheme provided efficiently computes $\mathrm{op}_{n,n-1}(123)$, which is sequence A220101 in the Online Encyclopeida of Integer Sequences [16].

Again, *Findrec* predicts the linear recurrence:

$$\mathrm{op}_{n,n-1}(123) = \begin{cases} 1 & n = 2 \\ \dfrac{(4n-6)(n-1)^2}{(n-2)^2(n+1)} \mathrm{op}_{n-1,n-2}(123) & n > 2. \end{cases}$$

While it remains an open problem to prove this recurrence describes our sequence in general, the asymptotic behavior of the sequence resulting from this recurrence is

$$\sim 2 \cdot 4^n \cdot \frac{\left(1 - \frac{25}{8n}\right)}{\sqrt{n}}.$$

We also use our recurrence to make the following conjecture, verified up to $n = 37$:

Conjecture 1.

$$op_{n,n-1}(123) = \frac{6(n-1)^2}{(n+1)!} \prod_{i=3}^{n} (4i-6)$$

The above discussion certainly suggests that $\mathrm{op}_{n,k}(123)$ is not monotone in k; for example, for $n = 4$ the sequence $\{\mathrm{op}_{n,k}(123) : 1 \leq k \leq 4\}$ is $\{1, 14, 27, 14\}$, as seen by elementary counting, the data above, and the fact that $C_4 = 14$. In a similar vein, if Conjecture 1 is true, it would show that $\mathrm{op}_{n,n-1}(123) \sim K \cdot 4^n/\sqrt{n}$, whereas it is well known that $\mathrm{op}_{n,n}(123) = C_n \sim K' 4^n/n^{3/2}$, a smaller number. We provide below a first result aimed at understanding the monotonicity (or lack thereof) discussed above:

Theorem 8. *$op_{n,4}(123) > op_{n,3}(123)$ for n sufficiently large.*

Proof. We start with a partial injection. Consider a partition p with three blocks $B_1/B_2/B_3$ with no block being a singleton. If $n \in B_3$ we map p to $p' := B_1/B_2/\{n\}/B_3 \setminus \{n\}$. If $n \in B_2$, we map p to $B_1/\{n\}/B_2 \setminus \{n\}/B_3$, and if $n \in B_1$ we map p to the 4-partition $p' = \{n\}/B_1 \setminus \{n\}/B_2/B_3$. It is clear that each of the 4-partitions avoid 123 and that the mapping $p \hookrightarrow p'$ is an injection. Thus the number of 3-avoiders with each block having 2 or more elements is less than the number of 4 avoiders with one block being $\{n\}$ and up to one more singleton block.

Simple over-counting shows that the number of 3-partitions (not necessarily avoiding) with at least one singleton block is no more than $3 \cdot n \cdot 2^{n-1}$ (we choose the singleton in n ways; place it in one of 3 positions; and then choose the two other parts by selecting any subset of the remaining elements). It is now left to show that the number of 4-avoiders with no singleton block is at least as large as $3 \cdot n \cdot 2^{n-1}$. We construct a lower bound on these by the following three-step process:

i make a 123 avoiding 3-partition of $\{3, 4, ..., n-3\}$ in roughly $\frac{(n-5)^2}{8} \cdot 2^{n-5}$ ways (the exact number is given by Theorem 2);

ii add $n-2$, 2 and 1 to the first, second and third blocks respectively;

iii define the first block to be $\{n-1, n\}$.

This completes the proof since $3 \cdot n \cdot 2^{n-1} < \frac{(n-5)^2}{8} \cdot 2^{n-5}$ for n sufficiently large.

□

We end with one more conjecture that states that monotonicity holds in a certain restricted sense.

Conjecture 2. *For each fixed k, there exists $n_0(k)$ such that for each $\rho \in S_m$ and $n \geq n_0(k)$,*

$$\mathrm{op}_{n,k+1}(\rho) > \mathrm{op}_{n,k}(\rho) > \cdots > \mathrm{op}_{n,|\rho|}(\rho).$$

7 A Stanley-Wilf Type Result

The Stanley-Wilf conjecture states that for every permutation $\rho \in \mathcal{S}_m$, there is a constant C such that the number $|S_n(\rho)|$ of permutations of length n which avoid ρ is asymptotic to C^n. The conjecture was first proved by Marcus and Tardos [12]. We will prove a similar result for ordered partitions.

Let $\rho \in \mathcal{S}_m$. Define $\mathcal{OP}*_{n,k}(\rho)$ be the set of ordered ρ-avoiding partitions of $[n]$ with k blocks, where some of the blocks may be empty. Let $\mathrm{op}*_{n,k}(\rho) = |\mathcal{OP}*_{n,k}(\rho)|$. We first prove, using Fekete's 1923 lemma for subadditive functions (see [18]), that a Stanley-Wilf [12] type result holds for $\mathrm{op}*_{n,k}(\rho)$.

Theorem 9. $\lim_{n\to\infty}$ *$op*^{1/n}_{n,k}(\rho)$ exists as a real number in $[1, \infty)$ for each fixed k and ρ.*

Proof. Let $\rho \in \mathcal{S}_\ell$ with $\ell \leq k$. Fix m, n, and consider $\pi \in \mathcal{OP}*_{m+n,k}(\rho)$. We shall show that π uniquely determines a pair (π_1, π_2) where $\pi_1 \in \mathcal{OQ}_{m,k}(\rho)$ and $\pi_2 \in \mathcal{OP}*_{n,k}(\rho)$. The mapping $\pi \hookrightarrow (\pi_1, \pi_2)$ that does this is the one that defines π_1 as the original partition with only the numbers $\{1, 2, \ldots, m\}$ placed in the same k blocks as before, and with π_2 defined as the original partition with only the numbers $\{m+1, m+2, \ldots, m+n\}$ placed again in the same k blocks as before, but renumbered as $\{1, 2, \ldots, n\}$. To give an example, with $m = 5; n = 6$, the ordered 321-avoiding block partition $1/\emptyset/3, 4, 6, 10/2, 5/7, 8/9, 11$ decomposes into the two 321-avoiding parts $\pi_1 = 1/\emptyset/3, 4/2, 5/\emptyset/\emptyset$ and $\pi_2 = \emptyset/\emptyset/1, 5/\emptyset/2, 3/4, 6$. It follows that

$$\mathrm{op}*_{m+n,k}(\rho) \leq \mathrm{op}*_{m,k}(\rho) \cdot \mathrm{op}*_{n,k}(\rho),$$

or that

$$\log \mathrm{op}*_{m+n,k}(\rho) \leq \log \mathrm{op}*_{m,k}(\rho) + \log \mathrm{op}*_{n,k}(\rho),$$

which shows that the function $\log \mathrm{op}*_{n,k}(\rho)$ is subadditive. Fekete's lemma ([18]) thus shows that

$$\lim_{n\to\infty} \frac{\log \mathrm{op}*_{n,k}(\rho)}{n} = \inf \frac{\log \mathrm{op}*_{n,k}(\rho)}{n};$$

in particular $\lim_{n\to\infty} \frac{\log \mathrm{op}*_{n,k}(\rho)}{n}$ *exists* as a number in $I = [-\infty, \infty)$ since the infimum of a non-empty real sequence is always in the interval I. Since in our case $\mathrm{op}*_{n,k}(\rho) \geq 1$, it follows that $\lim_{n\to\infty} \frac{\log \mathrm{op}*_{n,k}(\rho)}{n} \in [0, \infty)$, and thus $\lim_{n\to\infty} \mathrm{op}*^{1/n}_{n,k}(\rho) \in [1, \infty)$, as claimed. ☐

We now show that a Stanley-Wilf limit exists even when blocks are not allowed to be empty, and that the corresponding limit is the same as that for $\mathrm{op}*_{n,k}(\rho)$.

Theorem 10. $\lim_{n\to\infty} op^{1/n}_{n,k}(\rho)$ *exists in* $[1, \infty)$ *for each fixed* k *and* ρ.

Proof. Let $\rho \in S_m$. Suppose that $m \leq k$. We will prove the result by showing that there is a function $\phi(k)$ that satisfies $\frac{\mathrm{op}*_{n,k}(\rho)}{\phi(k)} \leq \mathrm{op}_{n,k}(\rho) \leq \mathrm{op}*_{n,k}(\rho)$, and $\lim_{n\to\infty} \phi(k)^{1/n} = 1$.

The second inequality is trivial. Assume that $k \leq n$. We prove the first inequality by providing an injection from $\mathcal{OP}*_{n,k}(\rho)$ into $\mathcal{OP}_{n,k}(\rho) \times [0, k]^{2k}$, where $[0, k]^{2k}$ is the number of words with $2k$ letters using the alphabet $[0, k] = \{0, 1, \ldots, k\}$.

Let $\pi \in \mathcal{OP}*_{n,k}(\rho)$, and assume that ρ does not end with m. We construct $(\hat{\pi}, w) \in \mathcal{OP}_{n,k} \times [0, k]^{2k}$ in the following way. First, move all of the empty blocks of π to the end of π keeping the relative order of the nonempty blocks unchanged, and call this new partition π_0. (Note that if ρ ends with m we move the empty blocks to the beginning of π and proceed as below.) Now, the first k letters of w are given by $w_i = j$ if the i^{th} block of π_0 was the j^{th} block of π, and $w_i = 0$ if the i^{th} block of π_0 is empty. Since the relative order of the nonempty blocks of π have not changed π_0 must be ρ avoiding.

Now, suppose there are a_1 empty blocks in π_0. Remove the a_1 largest elements of π_0 (that is the elements $n - a_1 + 1$ through n), and put one in each of the empty blocks at the end of π_0 so that they are in increasing order. Call this new partition π_1.

There can be no copies of ρ involving only the first $k - a_1$ blocks since such a copy would have been a copy in π_0 as well. If a copy of ρ involves one of the last a_1 blocks, then the element in the last block used would have to represent m since it would be the largest of all of the elements from π_1 used. This is impossible, since ρ does not end in m.

Suppose there are a_2 empty blocks in π_1. If $a_2 = 0$ then we are done, and we let $\pi_1 = \hat{\pi}$. If $a_2 > 0$ then we take the elements $n - a_1 - a_2 + 1$ through $n - a_1$ out of their blocks and place them in the a_2 empty blocks in such a way that the sequence formed by their placement is order isomorphic to the sequence created by the maxima of these blocks in π_0. Call this new partition π_2.

As before there can be no copies of ρ involving any of the last a_1 blocks of π_2. Thus, the only possible copies of ρ could be in the first $n - a_1$ blocks of π_2, and such a copy must involve at least one of the singleton blocks that was an empty block in π_1. Such a copy of ρ cannot only involve these formerly empty blocks of π_1 since this would imply a copy of ρ in π_0. Thus, a copy of ρ must involve elements from an original block from π_1 and some of the formerly empty blocks.

Suppose such a copy exists, call it $\sigma = \sigma_1\sigma_2 \cdots \sigma_m$. Suppose that $\sigma_{i_1}\sigma_{i_2} \ldots \sigma_{i_t}$ is the subsequence of elements from the formerly empty blocks. Replace each of these entries in σ by the maxima of the corresponding blocks in π_0. Call this new permutation τ. Now, τ is order isomorphic to σ since the sequence $\sigma_{i_1}\sigma_{i_2} \ldots \sigma_{i_t}$ is order isomorphic to the sequence of maxima of the corresponding blocks in π_0, and $\min\{\sigma_{i_j} : 1 \leq j \leq t\} > \max\{\sigma_s : s \notin \{i_j : 1 \leq j \leq t\}\}$. Thus, τ is a copy of ρ in π_0, which contradicts the fact that π_0 avoids ρ.

Suppose we reach a partition π_i with a_{i+1} empty blocks. We remove the largest a_{i+1} that are in blocks that have not been empty at any point during the construction. We fill the empty blocks with these a_{i+1} elements by placing one in each block so that they are order isomorphic to the maxima of the corresponding blocks in π_{i-1}. Thus forming π_{i+1}.

We must show that π_{i+1} avoids ρ. Suppose $\sigma = \sigma_1\sigma_2 \cdots \sigma_m$ is a copy of ρ in π_{i+1}. We know that none of the last a_1 blocks are involved. Form a permutation $\tau = \tau_1\tau_2 \ldots \tau_m$ from σ in the following way. If σ_i is in a block that has not yet been emptied then $\tau_i = \sigma_i$. If σ_i is in a block that was empty in π_j then τ_i is the maximum of the corresponding block in π_{j-1}. Now, τ is order isomorphic to σ by an argument similar to that for π_2. Thus, τ is a copy of ρ in π_0.

Once we reach π_j with no empty blocks, we set $\pi_j = \hat{\pi}$. Such a π_j will always be obtained since we assumed that $n \geq k$.

The final k letters of the word w are as follows. Let $w_{k+i} = 0$ if the i^{th} block of $\hat{\pi}$ has more than one element. Let $w_{k+i} = j$ if the i^{th} block has one element and that element was in the j^{th} block of π_0. The nature of w makes this process invertible and hence we have an injection. Since $|[0,k]^{2k}| = (1+k)^{2k}$, we have that $\frac{\mathrm{op^*}_{n,k}(\rho)}{\phi(k)} \leq \mathrm{op}_{n,k}(\rho)$ where $\phi(k) = (1+k)^{2k}$. □

An example will certainly help make the previous construction easier to understand. Suppose that $\rho = 132$. We have that $\pi = 8/\emptyset/359/12/\emptyset/46/7 \in \mathcal{OQ}_{9,7}(\rho)$. In the first step we move the empty blocks to the end, and obtain $\pi_0 = 8/359/12/46/7/\emptyset/\emptyset$. The first seven letters of the word w are 1346700 since the last 2 blocks were empty, the second block in π_0 was the third block in π, etc.

Now, we remove 8 and 9 from π_0 and use them to fill in the empty blocks at the end by placing them in increasing order to obtain, $\pi_1 = \emptyset/35/12/46/7/8/9$. The first block has been emptied by removing the 8.

We remove 7 from its block in π_1 and place it in the empty block. Notice that since there is only one element to place we do not need to worry about placing it in an order isomorphic way. This gives us $\pi_2 = 7/35/12/46/\emptyset/8/9$. The fifth block has been emptied by removing the 7.

We remove 6 from its block in π_2 and place it in the empty block obtaining $\pi_3 = 7/34/12/4/6/8/9$. No blocks of π_3 are empty, so we set $\hat{\pi} = \pi_3$. The last seven letters of the word w are 5004412. The second and third of these last seven letters are zeros since the second and third blocks of $\hat{\pi}$ have at least two elements each. The first letter is 5 since 7 was in the fifth block of π_0 etc.

This gives us the pair $(7/35/12/4/6/8/9, 13467005004412) \in \mathcal{OP}_{9,7}(132) \times [0,k]^{2k}$.

The proof of Theorem 10 can be substantially simplified if the monotonicity conjecture 2 is proved, since we would have that $\mathrm{oq}(n,k) \leq \sum_{j=1}^{k} \binom{k}{j} \mathrm{op}(n,j)$, where j indicates which of the k blocks are to be non-empty; varying the positions of these yields all possibilities for empty ordered blocks. Thus by monotonicity,

$$\mathrm{oq}(n,k) \leq (2^k - 1) \max_{1 \leq j \leq k} \mathrm{op}(n,j) = (2^k - 1)\mathrm{op}(n,k),$$

for n sufficiently large, which proves the first inequality in Theorem 10 with $\phi(k) = 2^k - 1$.

8 Future work/Open questions

We propose the following open questions for study.

i Enumeration of classes of avoidance numbers $\mathrm{op}_{n,k}(\rho)$, $k \geq 4$, is certainly of critical importance, starting with the case $|\rho| = 3$;

ii We know that $\lim \mathrm{op}_{n,3}^{1/n}(123) = 2$, and $\lim \mathrm{op}_{n,n}^{1/n}(123) = 4$. For which k does, e.g., $\lim \mathrm{op}_{n,k}^{1/n}(123) = 3$? Is $\sup_{k=k_n} \lim \mathrm{op}_{n,k}^{1/n}(123) < \infty$? Is $\sup_{k=k_n} \lim \mathrm{op}_{n,k}^{1/n}(123) = 4$?

iii To what extent are Conjectures 1 and 2 true?

iv Are the numbers $\mathrm{op}_{n,k}(\rho)$ unimodal for $3 \leq k \leq n$ and where do they attain their maximum?

References

[1] A. Burstein, *Enumeration of words with forbidden patterns*, Ph.D. thesis, University of Pennsylvania, 1998.

[2] D. Callan, A recursive bijective approach to counting permutations containing 3-letter patterns, Preprint **arXiv:math/0211380**.

[3] A. Goyt, Avoidance of partitions of a three element set, *Advances in Applied Mathematics* **41** (2008) 95–114.

[4] A. Goyt and B. Sagan, Set Partition Statistics and q-Fibonacci Numbers, *European Journal of Combinatorics* **30** (2009) 230–245.

[5] A. Goyt and L. Pudwell, Avoiding colored partitions of lengths two and three, *Pure Mathematics and Applications* **22**(2) (2011), 107–128.

[6] A. Goyt and L. Pudwell, Avoiding colored partitions of two elements in the pattern sense, *Journal of Integer Sequences* **15** (2012), 12.6.2.

[7] V. Jelínek and T. Mansour, On pattern-avoiding partitions, *Electron. J. Combin.*, **15(1):#R39** (2008)

[8] M. Klazar, On abab-free and abba-free set partitions, *European J. Combin.* **17** (1996), 53–68.

[9] M. Klazar, Counting pattern-free set partitions. I. A generalization of Stirling numbers of the second kind, *European J. Combin.* **21** (2000), 367–378.

[10] M. Klazar, Counting pattern-free set partitions. II. Noncrossing and other hypergraphs, *Electron. J. Combin.* **7** (2000) 25pp (electronic).

[11] D. E. Knuth, The art of computer programming. Volume 3. Sorting and Searching, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1973.

[12] A. Marcus and G. Tardos, Excluded permutation matrices and the Stanley-Wilf conjecture, *J. Combin. Theory Ser. A*, **107** (2004), no. 1, pp. 153–160.

[13] L. Pudwell, Enumeration schemes for words avoiding permutations. In *Permutation Patterns (2010)*, S. Linton, N. Ruskuc, and V. Vatter, Eds., vol. 376 of London Mathematical Society Lecture Note Series, Cambridge University Press, pp. 193–211. Cambridge: Cambridge University Press.

[14] B. Sagan, Pattern Avoidance in Set Partitions, *Ars Combin.*, **94** (2010), 79–96.

[15] R. Simion and F. W. Schmidt, Restricted permutations, *European J. Combin.* **6** (1985), no. 4, 383–406.

[16] N. Sloane, The Encyclopedia of Integer Sequences. Available at `http://oeis.org`, 2012.

[17] R. P. Stanley, *Enumerative Combinatorics, Vol. 2*, Cambridge University Press, 1999.

[18] J. Michael Steele, *Probability Theory and Combinatorial Optimization,* SIAM, Philadelphia, 1997.XC

[19] V. Vatter, Enumeration schemes for restricted permutations, *Combinatorics, Probability and Computing* **17** (2008), 137–159.

[20] J. Wimp and D. Zeilberger, Resurrecting the asymptotics of linear recurrences, *J. Math. Anal. Appl.* **111** (1985), 162–177.

[21] D. Zeilberger, Enumeration Schemes, and More Importantly, Their Automatic Generation, *Annals of Combinatorics* **2** (1998), 185–195.

[22] D. Zeilberger, FindRec: A Maple package that guesses recurrence equations for discrete functions of one variable, Available at `http://www.math.rutgers.edu/~zeilberg/EM08/FindRec.txt`, 2008.

[23] D. Zeilberger, AsyRec: A Maple package for Computing the Asymptotics of Solutions of Linear Recurrence Equations with Polynomial Coefficients, Available at `http://www.math.rutgers.edu/~zeilberg/tokhniot/AsyRec`, 2008.